



MEETING ABSTRACT

Open Access

Mechanical variations induce by foot orthoses on calcaneal fracture

T Breard^{1,2,3*}, M Janin^{1,2}

From 4th Congress of the International Foot and Ankle Biomechanics (i-FAB) Community
Busan, Korea. 8-11 April 2014


This work is meant to quantify the benefits of the foot orthoses [1,2] through the clinical case of a female patient aged 67 who broke her right calcaneum.

To investigate this, we implemented three experiments commonly used during podiatric examinations to assess walking parameters: the passive antepulsion test, the stabi-baro-podometrie analysis [3,4] and the Latero-Medial Index, measures taken immediately (T0) and after 16 days (T16) wearing plantar orthotics. (figure 1) Foot orthoses, deduced after clinical examination and quantitative analysis

of walking, are molded, which are supplemented by the addition of specific low stimulations. The results clearly show the benefits on stability. The foot orthoses allows the patient to recover the normal use of the ankle thanks to the positive effects on support and movement of a fractured foot [5]. Therefore, the foot orthoses tends to improve the balance of the fractured foot. Moreover, these positive effects are persistent throughout the time.

In a latter phase, the adjustments carried out on the foot orthoses that modify the foot simulation [3,4],

T0, Force Cuves without foot orthose. left / right



T16, Force Cuves without foot orthoses. left / right




Figure 1 T0, Force Cuves without foot orthose. Left/Right and T16, Force Cuves without foot orthoses. Left/right

¹Applied Podiatry College, 7 Treguel, 86000 Poitiers, France
Full list of author information is available at the end of the article

result in the improvement of the assessed parameters (static and dynamic). These variations tend to prove the benefits of the foot orthoses and justify the podiatric approach developed on this clinical case.

Authors' details

¹Applied Podiatry College, 7 Treguel, 86000 Poitiers, France. ²Podiatrist, PhD, Clinic, 7 Treguel, 86000 Poitiers, France. ³Maison médical de Roaillan, 33210 Roaillan, France.

Published: 8 April 2014

References

1. Murley GS, Landoir KB, Menz HB: Do foot orthoses change lower limb muscle activity in flat-arched feet towards a pattern observed in normal-arched feet? *Clin Biomech* 2010, **25**:728-736.
2. Murley GS, Landoir KB, Menz HB, Bird AR: Effect of foot posture, foot orthoses and footwear on lower limb muscle activity during walking and running: a systematic review. *Gait Posture* 2009, **29**:172-87.
3. McPoil TG, Cornwall MW: Use of plantar contact area to predict medial longitudinal arch height during walking. *JAMPA* 2006, **96**:489-494.
4. Wong L, Hunt A, Burns J, Crosbie J: Effect of foot morphology on center-of-pressure excursion during barefoot walking. *JAPMA* 2008, **98**:112-127.
5. Gillespie WJ, Gillespie LD, Paker MJ: Hip protectors for preventing hip fractures in older people. *Cochrane Database Syst. Rev* 2010, **6**(10): CD001255.
6. Janin M, Dupui P: The effects of unilateral medial arch support stimulation on plantar pressure and center of pressure adjustment in young gymnasts. *Neuroscience Letters* 2009, **461**:245-248.
7. Janin M, Toussaint L: Change in center of pressure with stimulations via anterior orthotic devices. *Gait Posture* 2005, **21**:1.579.

doi:10.1186/1757-1146-7-S1-A9

Cite this article as: Breard and Janin: Mechanical variations induce by foot orthoses on calcaneal fracture. *Journal of Foot and Ankle Research* 2014 **7**(Suppl 1):A9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

